

PRINCIPALLY SPECIALIZED CHARACTERS OF $\widehat{\mathfrak{sl}}(m|1)$ -MODULES

TAKUYA MURAKAMI

ABSTRACT. In this paper, we calculate a series of principally specialized characters of the $\widehat{\mathfrak{sl}}(m|1)$ -modules of level 1. In particular, we show that the principally specialized characters of the basic modules $L(\Lambda_0)$ is expressed as an infinite product. In addition, we deduce the specialized character formula of “quasiparticle” type.

0. INTRODUCTION

Character formulas of basic representation of $\widehat{\mathfrak{sl}}(m|n)$ are given in [KW2], based on their explicit construction in terms of bosonic and fermionic fields.

In this paper, we calculate the principally specialized characters of some $\widehat{\mathfrak{sl}}(m|1)$ -modules. In §1, we describe that the principally specialized characters of the basic $\widehat{\mathfrak{sl}}(m|1)$ -modules $L(\Lambda_0)$ is expressed as an infinite product. In §2, we deduce the specialized character formula of “quasiparticle” type.

We follow notation and terminologies from [KW2] without repeating their explanation.

1. SPECIALIZED CHARACTER FORMULA FOR SOME SERIES OF $\widehat{\mathfrak{sl}}(m|1)$ -MODULES

Throughout this paper, we assume that $m \geq 2$ and let $\alpha_0, \dots, \alpha_m$ denote the set of simple roots for $\widehat{\mathfrak{sl}}(m|1)$ where α_0 and α_m are odd and $\alpha_i (i = 1, \dots, m-1)$ are even. Provided that all s_i are positive integers, the sequence $s = (s_0, \dots, s_m)$ defines a homomorphism $\mathcal{F}_s : \mathbb{C}[[e^{-\alpha_0}, \dots, e^{-\alpha_m}]] \longrightarrow \mathbb{C}[[q]]$ by $\mathcal{F}_s(e^{-\alpha_i}) = q^{s_i} (i = 0, \dots, m)$, called the *specialization of type s* . In this paper, we consider the specialization of type $s = (1, \dots, 1, 0)$ which makes sense for the characters of integrable representations, and we write simply $\mathcal{F}$ for this specialization $\mathcal{F}_s$ when no confusion can arise.

Since the set of simple roots for the even part of $\widehat{\mathfrak{gl}}(m|1)$ is

$$\widehat{\Pi}' = \{\alpha'_0 = \alpha_0 + \alpha_m, \alpha_1, \dots, \alpha_{m-1}\},$$

this specialization is the principal specialization with respect to the even part of $\widehat{\mathfrak{gl}}(m|1)$; namely

$$\mathcal{F}(e^{-\alpha'_0}) = \mathcal{F}(e^{-\alpha_1}) = \dots = \mathcal{F}(e^{-\alpha_{m-1}}) = q.$$

We recall the Fock space and its charge decomposition:

$$F = \bigoplus_{s \in \mathbb{Z}} F_s$$

from §3 of [KW2].

Date: February 7, 2008.

Lemma 1.1. *Let $m \geq 2, s \in \mathbb{Z}$. Then we have:*

$$(a) \quad q^{\frac{sm}{2}} \mathcal{F}(e^{-\Lambda_0} chF_s) + q^{-\frac{sm}{2}} \mathcal{F}(e^{-\Lambda_0} chF_{-s}) = \frac{2 \prod_{i=1}^{\infty} (1+q^i)^2}{\varphi(q^m)^2}.$$

$$(b) \quad \mathcal{F}(e^{-\Lambda_0} chF_s) = \mathcal{F}(e^{-\Lambda_0} chF_{m-1-s}).$$

Here and further $\varphi(q) = \prod_{j=1}^{\infty} (1 - q^j)$.

Proof. In the case $\widehat{\mathfrak{gl}}(m|1)$, the formula (3.15) in [KW2] gives the following:

$$chF = e^{\Lambda_0} \prod_{k=1}^{\infty} \frac{\prod_{i=1}^m (1 + ze^{\epsilon_i - (k - \frac{1}{2})\delta})(1 + z^{-1}e^{-\epsilon_i - (k - \frac{1}{2})\delta})}{(1 - ze^{\epsilon_{m+1} - (k - \frac{1}{2})\delta})(1 - z^{-1}e^{-\epsilon_{m+1} - (k - \frac{1}{2})\delta})}, \quad (1.1)$$

where z is the “charge” variable.

Since $\alpha_0 = \delta - \epsilon_1 + \epsilon_{m+1}$ and $\alpha_i = \epsilon_i - \epsilon_{i+1}$, ($i = 1, \dots, m$), our principal specialization $\mathcal{F} = \mathcal{F}_s$ is written in terms of $e^{-\epsilon_i}$ as follows:

$$\mathcal{F}(e^{-\epsilon_i}) = q^{m-i} (i = 1, \dots, m), \quad \mathcal{F}(e^{-\epsilon_{m+1}}) = 1. \quad (1.2)$$

Thus, we obtain

$$\begin{aligned} \mathcal{F}(e^{-\Lambda_0} chF) &= \prod_{k=1}^{\infty} \frac{\prod_{i=1}^m (1 + zq^{i-m}q^{m(k-\frac{1}{2})})(1 + z^{-1}q^{m-i}q^{m(k-\frac{1}{2})})}{(1 - zq^{m(k-\frac{1}{2})})(1 - z^{-1}q^{m(k-\frac{1}{2})})} \\ &= \prod_{k=1}^{\infty} \frac{(1 + zq^{k-\frac{m}{2}})(1 + z^{-1}q^{k-1+\frac{m}{2}})}{(1 - zq^{m(k-\frac{1}{2})})(1 - z^{-1}q^{m(k-\frac{1}{2})})} \\ &= \prod_{k=1}^{\infty} \frac{(1 + (zq^{\frac{1-m}{2}})q^{k-\frac{1}{2}})(1 + (z^{-1}q^{\frac{m-1}{2}})q^{k-\frac{1}{2}})}{(1 + (-z^{-1})q^{m(k-\frac{1}{2})})(1 + (-z)q^{m(k-\frac{1}{2})})}. \end{aligned} \quad (1.3)$$

In order to compute the coefficient of z^s , we use the Jacobi triple product identity:

$$\prod_{n=1}^{\infty} (1 + zq^{n-\frac{1}{2}})(1 + z^{-1}q^{n-\frac{1}{2}}) = \frac{1}{\varphi(q)} \sum_{j \in \mathbb{Z}} z^j q^{\frac{1}{2}j^2}, \quad (1.4)$$

and also the following well-known identity (see (5.26) in [KP] and §5.8 in [K4]):

$$\begin{aligned} \prod_{k=1}^{\infty} (1 + zq^{k-\frac{1}{2}})^{-1} (1 + z^{-1}q^{k-\frac{1}{2}})^{-1} &= \frac{1}{\varphi(q)^2} \sum_{m \in \mathbb{Z}} (-1)^m \frac{q^{\frac{1}{2}m(m+1)}}{1 + zq^{m+\frac{1}{2}}} \\ &= \varphi(q)^{-2} \left(\sum_{m,k \geq 0} - \sum_{m,k < 0} \right) ((-1)^{m+k} z^k q^{\frac{1}{2}m(m+1) + (m+\frac{1}{2})k}). \end{aligned} \quad (1.5)$$

Replacing z by $zq^{\frac{1-m}{2}}$ in (1.4), z by $-z^{-1}$ and q by q^m in (1.5), we rewrite the right side of (1.3) by

$$\mathcal{F}(e^{-\Lambda_0} chF) = \frac{1}{\varphi(q)\varphi(q^m)^2} \sum_{k \in \mathbb{Z}} \left(\sum_{a,p \geq 0} - \sum_{a,p < 0} \right) (-1)^a z^{k-p} q^{\frac{1}{2}k(k+1) - \frac{km}{2} + \frac{m}{2}a(a+1) + m(a+\frac{1}{2})p}$$

and thus we have

$$\mathcal{F}(e^{-\Lambda_0} chF_s) = \frac{1}{\varphi(q)\varphi(q^m)^2} \left(\sum_{a,p \geq 0} - \sum_{a,p < 0} \right) (-1)^a q^{\frac{1}{2}(p+s)(p+s+1) - \frac{sm}{2} + \frac{m}{2}a(a+1) + map}. \quad (1.6)$$

For convenience, we introduce the following functions:

$$\begin{aligned} f_s(a, p) &= (-1)^a q^{\frac{1}{2}(p+s)(p+s+1) - \frac{sm}{2} + \frac{m}{2}a(a+1) + map}, \\ h_s &= \left(\sum_{a,p \geq 0} - \sum_{a,p < 0} \right) f_s(a, p). \end{aligned} \quad (1.7)$$

To prove this lemma, it is sufficient to prove the following two equations:

$$\begin{aligned} \text{(a')} \quad & q^{\frac{sm}{2}} h_s + q^{-\frac{sm}{2}} h_{-s} = 2\varphi(q) \prod_{i=1}^{\infty} (1 + q^i)^2. \\ \text{(b')} \quad & h_s = h_{m-1-s}. \end{aligned}$$

First we shall prove (a').

Since $f_s(-a, -p-1) = q^{-sm} f_{-s}(a, p)$, we have

$$\sum_{\substack{a > 0 \\ p \geq 0}} f_s(a, p) = \sum_{a,p < 0} f_s(-a, -p-1) = q^{-sm} \sum_{a,p < 0} f_{-s}(a, p).$$

Hence we obtain

$$\begin{aligned} h_s &= \sum_{\substack{a > 0 \\ p \geq 0}} f_s(a, p) + \sum_{p \geq 0} f_s(0, p) - \sum_{a,p < 0} f_s(a, p) \\ &= q^{-sm} \sum_{a,p < 0} f_{-s}(a, p) + \sum_{p \geq 0} f_s(0, p) - \sum_{a,p < 0} f_s(a, p). \end{aligned}$$

We then have

$$q^{\frac{1}{2}sm} h_s = q^{-\frac{1}{2}sm} \sum_{a,p < 0} f_{-s}(a, p) + q^{\frac{1}{2}sm} \sum_{p \geq 0} f_s(0, p) - q^{\frac{1}{2}sm} \sum_{a,p < 0} f_s(a, p). \quad (1.8)$$

A straightforward computation replacing s by $-s$ yields

$$q^{-\frac{1}{2}sm} h_{-s} = q^{\frac{1}{2}sm} \sum_{a,p < 0} f_s(a, p) + q^{-\frac{1}{2}sm} \sum_{p \geq 0} f_{-s}(0, p) - q^{-\frac{1}{2}sm} \sum_{a,p < 0} f_{-s}(a, p). \quad (1.9)$$

By (1.8) and (1.9), we obtain

$$q^{\frac{1}{2}sm} h_s + q^{-\frac{1}{2}sm} h_{-s} = 2 \sum_{p \geq 0} q^{\frac{1}{2}p(p+1)}. \quad (1.10)$$

It is known (see e.g. §5 in [KW1]) that the right side of (1.10) has the following product expansion:

$$\sum_{p \geq 0} q^{\frac{1}{2}p(p+1)} = \prod_{k \geq 1} \frac{1 - q^{2k}}{1 - q^{2k-1}} = \varphi(q) \prod_{i=1}^{\infty} (1 + q^i)^2,$$

and hence (a') follows.

Next we shall prove (b').

Since $f_s(-a-1, 0) = -f_s(a, 0)$, we have

$$\sum_{a \geq 0} f_s(a, 0) = \sum_{a < 0} f_s(-a-1, 0) = - \sum_{a < 0} f_s(a, 0).$$

Hence we obtain

$$\begin{aligned} h_s &= \sum_{\substack{a \geq 0 \\ p > 0}} f_s(a, p) + \sum_{a \geq 0} f_s(a, 0) - \sum_{a, p < 0} f_s(a, p) \\ &= \sum_{\substack{a \geq 0 \\ p > 0}} f_s(a, p) - \sum_{a < 0} f_s(a, 0) - \sum_{a, p < 0} f_s(a, p) \\ &= \left(\sum_{\substack{a \geq 0 \\ p > 0}} - \sum_{\substack{a < 0 \\ p \leq 0}} \right) f_s(a, p). \end{aligned}$$

Replacing a by $-(a+1)$ and p by $-p$, we have

$$\begin{aligned} h_s &= \left(\sum_{a, p < 0} - \sum_{a, p \geq 0} \right) f_s(-a-1, -p) \\ &= \left(\sum_{a, p < 0} - \sum_{a, p \geq 0} \right) (-1)^{a+1} q^{(-p+s)(-p+s+1) - \frac{sm}{2} + \frac{m}{2}a(a+1) + m(a+1)p} \\ &= \left(\sum_{a, p \geq 0} - \sum_{a, p < 0} \right) (-1)^a q^{\frac{1}{2}(p-s)(p-s-1) - \frac{sm}{2} + \frac{m}{2}a(a+1) + m(a+1)p}. \end{aligned} \quad (1.11)$$

Replacing s by $m-1-s$ in (1.11) and comparing it with (1.7), we get (b'). $\square$

Using Lemma 1.1 inductively, we have the following.

Proposition 1.2. *Let $m \geq 2$ and $k \in \mathbb{Z}_+$. Then we have*

$$\begin{aligned} \mathcal{F}(e^{-\Lambda_0} chF_{(k+1)(m-1)}) &= \mathcal{F}(e^{-\Lambda_0} chF_{-k(m-1)}) \\ &= q^{k \frac{m(m-1)}{2}} \left(\sum_{|j| \leq k} (-1)^{k-j} q^{(k^2-j^2) \frac{m(m-1)}{2}} \right) \frac{\prod_{i=1}^{\infty} (1+q^i)^2}{\varphi(q^m)^2}. \end{aligned}$$

Proof. By Lemma 1.1, we get the recurrence formula

$$\mathcal{F}(e^{-\Lambda_0} chF_{s+m-1}) = q^{\frac{sm}{2}} \left(\frac{2 \prod_{i=1}^{\infty} (1+q^i)^2}{\varphi(q^m)^2} - q^{\frac{sm}{2}} \mathcal{F}(e^{-\Lambda_0} chF_s) \right). \quad (1.12)$$

We now set for $k \in \mathbb{Z}$, $s = k(m-1)$, $x = q^{\frac{m(m-1)}{2}}$ and

$$H_k = \mathcal{F}(e^{-\Lambda_0} chF_{k(m-1)}) \left(\frac{\prod_{i=1}^{\infty} (1+q^i)^2}{\varphi(q^m)^2} \right)^{-1}.$$

For the proof of this proposition, it is sufficient to show that

$$H_{k+1} = x^k \left(\sum_{|j| \leq k} (-1)^{k-j} x^{k^2-j^2} \right), \quad k \in \mathbb{Z}_+. \quad (1.13)$$

We shall show (1.13) by induction on k . Using these notation, (1.12) is rewritten as

$$H_{k+1} = x^k(2 - x^k H_k).$$

Assume that (1.13) is true for $k - 1$. Then we obtain

$$\begin{aligned} H_{k+1} &= x^k \left\{ 2 - x^k \cdot x^{k-1} \left(\sum_{|j| \leq k-1} (-1)^{k-1-j} x^{(k-1)^2-j^2} \right) \right\} \\ &= x^k \left(2 + \sum_{|j| \leq k-1} (-1)^{k-j} x^{k^2-j^2} \right) \\ &= x^k \left(\sum_{|j| \leq k} (-1)^{k-j} x^{k^2-j^2} \right), \end{aligned}$$

proving the proposition. $\square$

We consider principally specialized characters of $\widehat{\mathfrak{sl}}(m|1)$ -modules $L(\Lambda_{(s)})$ where $\Lambda_{(s)}$ are defined in Remark 3.2 of [KW2], given by

$$\Lambda_{(s)} = \begin{cases} \Lambda_s & \text{if } 0 \leq s \leq m \\ -(s-m)\Lambda_0 + (1+s-m)\Lambda_m & \text{if } s \geq m \\ (1-s)\Lambda_0 + s\Lambda_m + s\delta & \text{if } s \leq 0. \end{cases} \quad (1.14)$$

Theorem 1.3. *Let $m \geq 2$. Then we have the following:*

(a) *For $\Lambda = \Lambda_0$ or Λ_{m-1} , we have*

$$\mathcal{F}(e^{-\Lambda} chL(\Lambda)) = \frac{\prod_{i=1}^{\infty} (1+q^i)^2}{\varphi(q^m)}. \quad (1.15)$$

(b) *For $\Lambda = \{k(m-1) + 1\}\Lambda_0 - k(m-1)\Lambda_m$ ($k \in \mathbb{Z}$), we have*

$$\mathcal{F}(e^{-\Lambda} chL(\Lambda)) = \left(\sum_{|j| \leq |k|} (-1)^{k-j} q^{(k^2-j^2)\frac{m(m-1)}{2}} \right) \frac{\prod_{i=1}^{\infty} (1+q^i)^2}{\varphi(q^m)}. \quad (1.16)$$

Proof. To prove (a), we let $k = 0$ in Proposition 1.2. Then, since $chF_s = \varphi(e^{-\delta})^{-1} chL(\Lambda_{(s)})$ and $\Lambda_{(0)} = \Lambda_0$, we get

$$\mathcal{F}(e^{-\Lambda_0} chL(\Lambda_0)) = \frac{\prod_{i=1}^{\infty} (1+q^i)^2}{\varphi(q^m)}.$$

Since $\Lambda_{(m-1)} = \Lambda_{m-1} = \Lambda_0 - \alpha_m$, we have

$$\mathcal{F}(e^{-\Lambda_{(m-1)}}) = \mathcal{F}(e^{-\Lambda_0}) \mathcal{F}(e^{\alpha_m}) = \mathcal{F}(e^{-\Lambda_0}),$$

and hence

$$\begin{aligned} \mathcal{F}(e^{-\Lambda_{m-1}} chL(\Lambda_{m-1})) &= \mathcal{F}(e^{-\Lambda_0} chL(\Lambda_{(m-1)})) \\ &= \mathcal{F}(e^{-\Lambda_0} chF_{m-1}) \varphi(q^m) \\ &= \frac{\prod_{i=1}^{\infty} (1+q^i)^2}{\varphi(q^m)} \end{aligned}$$

which is (a).

To prove (b), we let $\Lambda = \{k(m-1) + 1\}\Lambda_0 - k(m-1)\Lambda_m$ for $k \in \mathbb{Z}$. First we consider in the case $k \in \mathbb{Z}_+$, since $\Lambda_m = \Lambda_0 - \frac{1}{m-1}\alpha_1 - \cdots - \frac{m}{m-1}\alpha_m$ and by (1.14), it follows that

$$\mathcal{F}(e^{-\Lambda_{(-k(m-1))}}) = \mathcal{F}(e^{-\Lambda})q^{-km(m-1)} = \mathcal{F}(e^{-\Lambda_0})q^{-\frac{km(m-1)}{2}},$$

and hence we obtain

$$\begin{aligned} \mathcal{F}(e^{-\Lambda}chL(\Lambda)) &= \mathcal{F}(e^{-\Lambda_{(-k(m-1))}}chL(\Lambda_{(-k(m-1))})) \\ &= q^{-\frac{km(m-1)}{2}}\mathcal{F}(e^{-\Lambda_0}chL(\Lambda_{(-k(m-1))})) \\ &= \left(\sum_{|j| \leq k} (-1)^{k-j} q^{(k^2-j^2)\frac{m(m-1)}{2}} \right) \frac{\prod_{i=1}^{\infty} (1+q^i)^2}{\varphi(q^m)}. \end{aligned}$$

In the case $k < 0$, the same discussion as above yields

$$\mathcal{F}(e^{-\Lambda_{((-k+1)(m-1))}}) = \mathcal{F}(e^{-\Lambda}) = \mathcal{F}(e^{-\Lambda_0})q^{\frac{km(m-1)}{2}},$$

and hence we obtain

$$\begin{aligned} \mathcal{F}(e^{-\Lambda}chL(\Lambda)) &= \mathcal{F}(e^{-\Lambda_{((-k+1)(m-1))}}chL(\Lambda_{((-k+1)(m-1))})) \\ &= q^{\frac{km(m-1)}{2}}\mathcal{F}(e^{-\Lambda_0}chL(\Lambda_{((-k+1)(m-1))})) \\ &= \left(\sum_{|j| \leq -k} (-1)^{k-j} q^{(k^2-j^2)\frac{m(m-1)}{2}} \right) \frac{\prod_{i=1}^{\infty} (1+q^i)^2}{\varphi(q^m)} \end{aligned}$$

which is (b). □

Remark 1.4. This is an additional remark to Theorem 1.3.

We consider an asymptotic behavior of principally specialized characters $\mathcal{F}(e^{-\Lambda_0}chL(\Lambda_0))$.

We shall write $f(\tau) \underset{\tau \downarrow 0}{\sim} g(\tau)$ if $\lim_{\tau \downarrow 0} f(\tau)/g(\tau) = 1$, where $\tau \downarrow 0$ means that $\tau = iT$ ($T > 0$) and $T \rightarrow 0$.

It is known (see e.g. (13.13.5) in [K3]) that the asymptotic behavior of the Dedekind η -function

$$\eta(\tau) = q^{\frac{1}{24}}\varphi(q) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n) \quad (q = e^{2\pi i\tau})$$

is given by

$$\eta(\tau) \underset{\tau \downarrow 0}{\sim} (-i\tau)^{-\frac{1}{2}} e^{-\frac{\pi i}{12\tau}},$$

and so

$$\varphi(q) = q^{-\frac{1}{24}}\eta(\tau) \underset{\tau \downarrow 0}{\sim} (-i\tau)^{-\frac{1}{2}} e^{-\frac{\pi i}{12\tau}}.$$

From this assertion, we have

$$\prod_{n=1}^{\infty} (1 + q^n) = \frac{\varphi(q^2)}{\varphi(q)} \underset{\tau \downarrow 0}{\sim} \frac{1}{\sqrt{2}} e^{\frac{\pi i}{24\tau}}$$

and hence

$$\begin{aligned} \mathcal{F}(e^{-\Lambda_0} chL(\Lambda_0)) &\underset{\tau \downarrow 0}{\sim} \frac{\frac{1}{2} e^{\frac{\pi i}{12\tau}}}{(-im\tau)^{-\frac{1}{2}} e^{-\frac{\pi i}{12m\tau}}} \\ &= \frac{\sqrt{m}}{2} (-i\tau)^{\frac{1}{2}} e^{\frac{2\pi i}{\tau} \frac{m+1}{24m}}. \end{aligned}$$

By applying the Tauberian theorem (see [I] and Proposition 4.22 in [KP]), we have

$$a_n \underset{n \rightarrow \infty}{\sim} \frac{1}{8\sqrt{3}n} (m+1)^{\frac{1}{2}} e^{\pi \sqrt{\frac{2}{3}} \frac{m+1}{m} n}$$

where $\mathcal{F}(e^{-\Lambda_0} chL(\Lambda_0)) = \sum_{n=0}^{\infty} a_n q^n$.

Next, we consider for principally specialized characters of some series of level 1 integrable modules. By Theorem 1.3, we have for $\Lambda = \Lambda_{m-1}$ or $\{k(m-1) + 1\}\Lambda_0 - k(m-1)\Lambda_m$ ($k \in \mathbb{Z}$),

$$\begin{aligned} \mathcal{F}(e^{-\Lambda} chL(\Lambda)) &\underset{\tau \downarrow 0}{\sim} \mathcal{F}(e^{-\Lambda_0} chL(\Lambda_0)) \\ &\underset{\tau \downarrow 0}{\sim} \frac{\sqrt{m}}{2} (-i\tau)^{\frac{1}{2}} e^{\frac{2\pi i}{\tau} \frac{m+1}{24m}}. \end{aligned} \quad (1.17)$$

The formula (4.12) in [KW2] is the asymptotics of the basic specialization (the specialization of type $(1, 0, \dots, 0)$) of $\widehat{\mathfrak{sl}}(m|1)$ -module $L(\Lambda_{(s)})$. The formula which is obtained by replaced τ by $m\tau$ in (4.12) of [KW2] coincides with (1.17).

2. SPECIALIZED CHARACTER FORMULA OF “QUASIPARTICLE” TYPE

Proposition 2.1. *For $m \geq 2$ and $s \in \mathbb{Z}$, we have*

$$\mathcal{F}(e^{-\Lambda_0} chF_s) = q^{-\frac{sm}{2}} \sum_{\substack{a,b,c,d \in \mathbb{Z}_+ \\ a-b+c-d=s}} \frac{q^{\frac{a(a+1)}{2}} q^{\frac{b(b-1)}{2}} q^{cm}}{(q)_a (q)_b (q^m)_c (q^m)_d}.$$

Here and further $(q)_n = \prod_{j=1}^n (1 - q^j)$.

Proof. This formula is shown just by the same argument as that in the proof of the formula (3.14) in [KW2], by making use of a basis of F_s :

$$\begin{aligned} &(\psi_{-(j_1, 1 - \frac{1}{2})}^{(1)} \cdots \psi_{-(j_1, a_1 - \frac{1}{2})}^{(1)}) \cdots (\psi_{-(j_m, 1 - \frac{1}{2})}^{(m)} \cdots \psi_{-(j_m, a_m - \frac{1}{2})}^{(m)}) \\ &\times (\psi_{-(j'_1, 1 - \frac{1}{2})}^{(1)*} \cdots \psi_{-(j'_1, b_1 - \frac{1}{2})}^{(1)*}) \cdots (\psi_{-(j'_m, 1 - \frac{1}{2})}^{(m)*} \cdots \psi_{-(j'_m, b_m - \frac{1}{2})}^{(m)*}) \\ &\times (\varphi_{-(k_1 - \frac{1}{2})}^{(1)} \cdots \varphi_{-(k_c - \frac{1}{2})}^{(1)}) (\varphi_{-(k'_1 - \frac{1}{2})}^{(1)*} \cdots \varphi_{-(k'_d - \frac{1}{2})}^{(1)*}) |0\rangle \end{aligned}$$

satisfying

$$\begin{aligned} 0 < j_{i,1} < \cdots < j_{i,a_i}, 0 < j'_{i,1} < \cdots < j'_{i,b_i} \quad (i = 1, \dots, m), \\ 0 < k_1 \leq \cdots \leq k_c, 0 < k'_1 \leq \cdots \leq k'_d \end{aligned}$$

and

$$a - b + c - d = s$$

where $a = \sum_i a_i, b = \sum_i b_i$.

Since

$$\begin{aligned} \text{weight}(\psi_k^{(i)}) &= \epsilon_i + k\delta, & \text{weight}(\psi_k^{(i)*}) &= -\epsilon_i + k\delta, \\ \text{weight}(\varphi_k^{(1)}) &= \epsilon_{m+1} + k\delta, & \text{weight}(\varphi_k^{(1)*}) &= -\epsilon_{m+1} + k\delta, \end{aligned}$$

and (1.2), we have

$$\begin{aligned} \mathcal{F}\left(e^{\text{weight}(\psi_{-(j-\frac{1}{2})}^{(i)})}\right) &= q^{i+m(j-\frac{3}{2})}, & \mathcal{F}\left(e^{\text{weight}(\psi_{-(j-\frac{1}{2})}^{(i)*})}\right) &= q^{-i+m(j+\frac{1}{2})}, \\ \mathcal{F}\left(e^{\text{weight}(\varphi_{-(j-\frac{1}{2})}^{(1)})}\right) &= q^{m(j-\frac{1}{2})}, & \mathcal{F}\left(e^{\text{weight}(\varphi_{-(j-\frac{1}{2})}^{(1)*})}\right) &= q^{m(j-\frac{1}{2})}, \end{aligned}$$

for $i = 1, \dots, m$ and $j \in \mathbb{N}$.

Hence we have

$$\begin{aligned} \mathcal{F}(e^{-\Lambda_0} ch F_s) &= \sum_{\substack{a,b,c,d \in \mathbb{Z}_+ \\ a-b+c-d=s}} \left(\sum_{0 < i_1 < \cdots < i_a} q^{(i_1 - \frac{m}{2}) + \cdots + (i_a - \frac{m}{2})} \right) \left(\sum_{0 < j_1 < \cdots < j_b} q^{(j_1 + \frac{m}{2} - 1) + \cdots + (j_b + \frac{m}{2} - 1)} \right) \\ &\quad \times \left(\sum_{0 < k_1 \leq \cdots \leq k_c} q^{(mk_1 - \frac{m}{2}) + \cdots + (mk_c - \frac{m}{2})} \right) \left(\sum_{0 < l_1 \leq \cdots \leq l_d} q^{(ml_1 - \frac{m}{2}) + \cdots + (ml_d - \frac{m}{2})} \right) \\ &= q^{-\frac{sm}{2}} \sum_{\substack{a,b,c,d \in \mathbb{Z}_+ \\ a-b+c-d=s}} \left(\sum_{0 < i_1 < \cdots < i_a} q^{i_1 + \cdots + i_a} \right) \left(q^{-b} \sum_{0 < j_1 < \cdots < j_b} q^{j_1 + \cdots + j_b} \right) \\ &\quad \times \left(\sum_{0 < k_1 \leq \cdots \leq k_c} q^{m(k_1 + \cdots + k_c)} \right) \left(q^{-dm} \sum_{0 < l_1 \leq \cdots \leq l_d} q^{m(l_1 + \cdots + l_d)} \right) \\ &= q^{-\frac{sm}{2}} \sum_{\substack{a,b,c,d \in \mathbb{Z}_+ \\ a-b+c-d=s}} \frac{q^{\frac{1}{2}a(a+1)}}{(q)_a} \frac{q^{\frac{1}{2}b(b-1)}}{(q)_b} \frac{q^{cm}}{(q^m)_c} \frac{1}{(q^m)_d}, \end{aligned}$$

proving the proposition. $\square$

By Theorem 1.3 and Proposition 2.1, we also have shown

Corollary 2.2. *For $m \geq 2$, the following formula holds*

$$\frac{\prod_{i=1}^{\infty} (1 + q^i)^2}{\varphi(q^m)^2} = \sum_{\substack{a,b,c,d \in \mathbb{Z}_+ \\ a-b+c-d=0}} \frac{q^{\frac{a(a+1)}{2}} q^{\frac{b(b-1)}{2}} q^{cm}}{(q)_a (q)_b (q^m)_c (q^m)_d}.$$

REFERENCES

- [HB] K. Hikami and B. Basu-Mallick, Supersymmetric Polychronakos spin chain: Motif, distribution function, and character, *Nuclear Physics B*, **566** (2000), 511–528.
- [I] A. E. Ingham, A Tauberian theorem for partitions, *Ann. of Math.*, **42** (1941), 1075–1090.
- [K1] V. G. Kac, Lie superalgebras, *Adv. Math.*, **26** (1977), 8–96.
- [K2] V. G. Kac, Infinite-dimensional algebras, Dedekind’s η -function, classical Möbius function and the very strange formula, *Adv. in Math.*, **30** (1978), 85–136.
- [K3] V. G. Kac, *Infinite dimensional Lie algebras*, 3rd edition, Cambridge University Press, 1990.
- [K4] V. G. Kac, *Vertex algebras for beginners*, University lecture series, Vol. **10**, AMS, Providence RI, 1996. Second edition, 1998.
- [KP] V. G. Kac and D. H. Peterson, Infinite-dimensional Lie algebras, theta functions and modular forms, *Adv. Math.*, **53** (1984), 125–264.
- [KW1] V. G. Kac and M. Wakimoto, Integrable highest weight modules over affine superalgebras and number theory, *Progress in Math.*, **123**, Birkhäuser, Boston, 1994, 415–456.
- [KW2] V. G. Kac and M. Wakimoto, Integrable highest weight modules over affine superalgebras and Appell’s function, to appear in *Commun. Math. Phys.*
- [S] R. P. Stanley, *Enumerative Combinatorics* Volume I, Wadsworth & Brooks / Cole Advanced Books & Software, 1986.

GRADUATE SCHOOL OF MATHEMATICS, KYUSHU UNIVERSITY 33, FUKUOKA, 812-8581, JAPAN
E-mail address: ma299040@math.kyushu-u.ac.jp